

ihr Stiel, dieser aber mehr als doppelt so lang als der Stiel der zweiten Hinterrandzelle; die erstere Gabelzelle beginnt erst etwas hinter der Discoidalzelle, bei der die kleine Querader auf der Mitte steht; die vierte Hinterrandzelle ist kurz gestielt. — 14 mm lang.

(Fortsetzung folgt.)

## A New Genus and Species of Aphrastobraconini from Borneo.

By P. Cameron.

### *Hewittella* gen. nov.

Eyes large, the malar space distinct. Mesonotum flat, the scutellum slightly raised above it. Metanotum with a depressed  $\wedge$ -shaped area in the centre; on the apical half is a stout keel bordered by 2 oblique ones on its lower half. Median cellule distinctly longer than the submedian. Otherwise as in *Jphiaulax*, with a long radial cellule. Temples moderately wide.

This genus differs from the type of the group (*Aphrastobracon* from Ceylon) in their being a distinct malar space and temples, and in their being a distinct area on the metanotum; in the former respect it departs from 2 of the characteristics of the tribe and agrees with *Odesia* Cam. from South Africa. The difference between it and *Odesia*\*) may be shown thus:

Radius not reaching to the apex of the wing, the radial cellule therefore short; mesonotum not flattened at the apex, 2<sup>nd</sup> cubital cellule narrowed in front; the 2<sup>nd</sup> abdominal segment longer than wide, the metanotum without an area . . . . . *Odesia*.

Radius reaching to the apex of the wing, the radial cellule long, mesonotum flattened at apex; 2<sup>nd</sup> cubital cellule long, not narrowed in front; the 2<sup>nd</sup> abdominal segment wider than long; the metanotum with a distinctly bordered area . . . . . *Hewittella*.

\*) *Odesia* Cam., Annals S. Af. Mus. V. 79.

**Hewittella flavomaculata** sp. nov. ♀.

Rufo-testaceous; the face, the greater part of the vertex, the orbits narrowly and more or less of the pleurae, mesonotum and scutellum, yellow; the antennae, tips of mandibles, 2 large quadrangular marks on the 5<sup>th</sup> abdominal segment and the hind legs, except the joints, black; wings hyaline, the base slightly suffused with yellow; the stigma and costa fuscous, the former pale at the base; the nervures are paler; the apical segments of the abdomen are pale testaceous. — Length 6 mm; terebra 1 mm.

Sarawak, Borneo (Hewitt).

Smooth, shining; a rugose space, narrowed above, over the mouth; the front depressed. Mesonotum and scutellum flat, especially the former at the apex; the latter is more elevated than the former. Abdomen closely rugosely punctured; the area on 2<sup>nd</sup> segment punctured, smooth at the base, its apex is produced shortly beyond the middle of the segment; this narrowed apical part is depressed in the centre, the sides sharp, forming keels; there are 3 distinctly defined, crenulated furrows; the apices of the segments are smooth. Legs covered with short pubescence. The amount of the black and yellow colour varies.

---

## Literatur.

**Adams, C. F.** *Notes on and descriptions of North American Diptera.* (Kansas Univ. Science Bull. Vol. II, No. 14. 1904, 433—455). Neue Arten: *Simulium notatum*, *Mydas abdominalis*, *Xylophagus nitidus* (Tabelle 435), *Chrysopila lucifera*, *bella*, *flavibarbis* (Tabelle für 36 Arten), *Symphoromyia flavipalpis* (439), *Leptis pleuralis*, *palpalis* (Tabelle 440), *Psilocephala occipitalis*, *lateralis* (444), *Thereva anomala*, *Scenopinus mirabilis*, *electa* (445), *Nausigaster scutellaris* (446), *Sphyximorpha Snowi* (447), *Tetanocera inopa*, *Ictericica fasciata* (Tabelle 449), *Urellia flava*, *conjuncta occidentalis* (Tabelle 450), *Elachiptera bilineata*, *Hippelates splendens* (453), *Limosina exigua*, *occidentalis*, *sordipes* (Tabelle 454).

---